



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{d^2 s}{dt^2} = -S \frac{dr}{ds} - S' \frac{dr'}{ds} \dots \dots (1).$$

But $S=m/r$, $S'=m/r'$, and (1) becomes

$$\frac{d^2 s}{dt^2} = -\frac{m}{r} \frac{dr}{ds} - \frac{m}{r'} \frac{dr'}{ds} \dots \dots (2).$$

Multiply by $2(ds/dt)$ and integrate ; then

$$\frac{ds^2}{dt^2} = -m \log r^2 - m \log r'^2 + C \dots \dots (3).$$

When $r=a$, $r'=a$, $\frac{ds}{dt}=\beta$; $\therefore C=\beta^2-m^2 \log \frac{1}{a^4}$, and (3) is

$$\beta^2 = m^2 \log \frac{1}{r^2 r'^2} + \beta^2 - m^2 \log \frac{1}{a^4} \dots (4),$$

or, $rr'=a^2 \dots \dots (5)$, a lemniscate.

[Other solutions of this problem will appear in the next issue.]

DIOPHANTINE ANALYSIS.

73. Proposed by M. A. GRUBER, A. M., War Department, Washington, D. C.

Find integral values for x and y in $\left(\begin{matrix} 2x^2 - y^2 = \square \\ 2y^2 - x^2 = \square \end{matrix} \right)$.

I. Solution by the PROPOSER.

$$2x^2 - y^2 = \square = a^2 \dots \dots (1), \quad 2y^2 - x^2 = \square = (2).$$

From (1), $y^2 = 2x^2 - a^2$. Substituting this in (2), we have $3x^2 - 2a^2 = b^2$. Whence $x = \frac{1}{3}\sqrt{3(2a^2 + b^2)}$, and $y = \frac{1}{3}\sqrt{3(a^2 + 2b^2)}$.

As far as I know, the only method of rationalizing both radicals is to put $a=b$. Then $x=y=a=b$.

Accordingly, no *different* integral values can be found for x and y .

This problem is the key to Problem 62, "To find four squares in arithmetical progression."

The roots of the four squares would then be, respectively,

$$a, \quad \frac{1}{3}\sqrt{3(2a^2 + b^2)}, \quad \frac{1}{3}\sqrt{3(a^2 + 2b^2)}, \quad b.$$

The common difference of the squares is $\frac{1}{9}(b^2 - a^2)$.

According to the above solution, the roots of the four squares could not all be rational integers ; *one* of them, at least, must be a *surd*. It is evident, however, that an infinite number of sets of four squares can be found in which *two* of the roots are rational integers.

Put $a=1$ and $b=2$. Then the roots of the four squares are 1, $\sqrt{2}$, $\sqrt{3}$, 2.

Put $a=1$ and $b=5$. Then the roots are 1, 3, $\sqrt{17}$, 5.

A similar proof was received from *CHARLES C. CROSS*.

II. Solution by *A. H. BELL*, Hillsboro, Ill.

Take $2y^2 - x^2 = \square$, or $x^2 - 2y^2 = -\square = -1 = -4$, etc.

In $x^2 - 2y^2 = -1 \dots (3)$, the integral values for x and y are the alternate convergent fractions for the $\sqrt{2}$ to

$$x/y = 1/1, 7/5, 41/29, \text{ etc.} \dots (4).$$

For the next, $x^2 - 2y^2 = -4$. $(4) \times 1/4$,

$$\frac{x}{y} = \frac{1 \times 2}{1 \times 2}, \quad \frac{7 \times 2}{5 \times 2}, \quad \frac{41 \times 2}{29 \times 2}, \quad \text{etc.}$$

Consequently the interchangeable values of x and y must be found in the first fraction and no other.

III. Solution by *JOSIAH H. DRUMMOND*, LL. D., Portland, Me.

$2x^2 - y^2 = \square \dots (1)$, and $2y^2 - x^2 = \square \dots (2)$.

Take $x=my$ and $2m^2 - 1 = \square \dots (3)$, and $2 - m^2 = \square \dots (4)$.

Then $m < \sqrt{2}$ and $m > \frac{1}{\sqrt{2}}$.

It is manifest that both (3) and (4) are rational when $m=1$, which is $< \sqrt{2}$ and $> \frac{1}{\sqrt{2}}$.

Then in (3) take $m=n+1$, and we have

$$2n^2 + 4n + 1 = \square = (\text{say})(qn-1)^2, \text{ whence}$$

$$n = \frac{2(q+2)}{q^2-2} \text{ and } m = n+1 = \frac{(q+1)^2+1}{q^2-2}.$$

Substituting this value of m in (4) and reducing by the usual methods, we find $q=0$ and $m=\pm 1$.

Hence $x=\pm y$ and the integral values are any equal numbers, positive or negative, or one positive and the other negative.

MISCELLANEOUS.

65. Proposed by *J. M. COLAW*, A. M., Monterey, Va.

Three circles, radii in ratio 1, 3, 5, are tangent externally and enclose one acre; what are the radii?